

Principle Outcomes of Concern

The following are the outcome measures drawn from the H2P2. Those outcome measures that are comparable to outcome measures found in *Healthy People 2010* are highlighted in bold.

– Infectious Diseases

- (IDO-1) **Incidence of Bacterial Meningitis**¹

– Cancer

- (CAO1.1-3) Reduce Incidence Rates of New Cancer by Site:
 - Lung
 - Breast
 - Prostate
- (CAO2.1-3) **Reduce Mortality Rates of New Cancer by Site**
 - **Lung**
 - **Breast**
 - **Prostate**
-
- (CAO3) Obtain regional data by Subarea Health Planning Regions
- (CAO4) Participation on [sic] Ongoing Cancer Research Activities

– Diabetes and Other Chronic Disabling Conditions

Diabetes

- (CDO-D1) **Diabetes Prevalence Rates**
 -
- (CDO-D2) Diabetic Related Inpatient Admission Rate
 -
- (CDO-D3) **Diabetic Limb Amputation Rate**
 -
- (CDO-D4) Ambulatory Care Sensitive (ACS) Admissions for Diabetes with Long Term Complications
 -
- (CDO-D5) **Diabetes-related Mortality Rate**
 -

Asthma

- (CDO-A1) Asthma Prevalence Rate
 -
- (CDO-A2) **Asthma-related Mortality Rate**
 -

Principle Outcomes of Concern

- (CDO-A3) **Asthma-related ER Visits**
—
- (CDO-A4) **Asthma-related Admissions Rate**
—
- (CDO-A5) Ambulatory Care Sensitive (ACS) Admissions for Pediatric Asthma
—
- (CDO-A6) Ambulatory Care Sensitive (ACS) Admissions for Adult Asthma
—
- (CDO-A7) Change in Population Severity
—
- (CDO-A8) Change in Anti-inflammatory to Beta-Agonist Ratio
—

Alzheimer's Disease

- (CDO-Z1) Alzheimer's prevalence rate
—
- (CDO-Z2) Alzheimer's mortality rate

— Heart Disease and Stroke

- (HDO1) **Proportion of people with high blood pressure who are receiving treatment for high blood pressure**
—
- (HDO2) Prevalence of elevated blood cholesterol and its components that exceed the recommended AHA standard
—
- (HDO3) Prevalence of overweight among people aged 20 and older and among adolescents aged 12 through 19
—
- (HDO4) Proportion of people aged 6 and older who engage regularly in light to moderate physical activity for at least 30 minutes per day
—
- (HDO5) Prevalence of cigarette smoking among people aged 20 and older
—
- (HDO6) **Deaths from CHD per 100,000 people in all ethnicities**
—
- (HDO7) **Stroke deaths per 100,000 people in all ethnicities in Hawaii**
—
- (HDO8) Open Heart procedures discharges
—
- (HDO9) Cardiac Cath discharges
—
- (HDO10) **Cardiac Interventions**
—
- (HDO11) Cardiac medicine discharges
—

- (HDO12) Other circulatory discharges
-
- (HDO13) Neurology medicine discharges
- *Preventable Injuries and Violence*
 - (PIO1) **Reduce childhood injury, death and hospitalization rates**
 - (PIO2) **Decrease rates of injury resulting from domestic violence**
 - (PIO3) Reduce alcohol- and drug-related injury rates
 - (PIO4) Reduce work-related injury and illness rate
 - (PIO5) **Reduce the rate of hip fracture hospitalizations among the elderly**
- *Dental (Oral) Health*
 - (DHO1.1) Decrease deaths from oral cancer
 - (DHO1.2) Reduce cigarette use
 - (DHO2) Reduce rates of sexually transmitted diseases
 - (DHO3.1) **Reduce decay incidence among 6 to 8 year olds**
 - (DHO3.2) **Reduce baby bottle caries among 5 year olds**
 - (DHO3.3) **Reduce dft (decayed teeth incidence)**
 - (DHO3.4) **Reduce decay to the general population percentage**
 - (DHO4) Decrease incidence of substance abuse
- *Maternal, Infant & Child Health*
 - Maternal Health*
 - (MHO1) **Mothers who received prenatal care in the first trimester of pregnancy**
 - (MHO2) Births to teenaged mothers age 15-19 years
 - (MHO3) **Pregnancies affected by hypertensive disorders**
 - (MHO4) **Pregnancies affected by infections**

Principle Outcomes of Concern

- (MHO5) Pregnancies affected by complications other than hypertensive disorders and infections
- (MHO6) **Premature labor and/or deliveries**

Infant Health

- (IHO1) Five-minute APGAR score of less than 7
- (IHO2) **Deaths of neonates under 28 days**
- (IHO3) **Deaths of infants weighing less than 2,500 g (under 28 days old)**

Pediatric Health

- (PHO1) Hospitalization for children 14 years of age and under for asthma
- (PHO2) Pediatric gastroenteritis hospital/ER admissions
- (PHO3) Number of pediatric patients seen in an ER for water related injuries
- (PHO4) Number of pediatric patients seen in an ER for burns
- (PHO5) Number of pediatric patients seen in an ER for trauma (excludes burns and water-related injuries)
- (PHO6) Number of pediatric patients who contracted rheumatic fever
- (PHO7) Number of pediatric patients seen in an ER for non-fatal poisonings
- (PHO8) Number of hospitalized cases due to child abuse and neglect
- (PHO9) **Childhood deaths**
- (PHO10) Motor vehicle death incidence of children 14 and under
- (PHO11) Deaths resulting from child abuse and neglect

Adolescent Health

- (AHO1) Teen Pregnancy (☐ 20 years of age)
- (AHO2) DSM-IV psychological disorders in the adolescent population
- (AHO3) Adolescents who used tobacco products in past month (ages 11-21)

- (AHO4) Adolescents who used alcohol in past month
- (AHO5) Adolescents who used marijuana in previous month
- (AHO6) Adolescents who used some type of illicit drug in previous month
- (AHO7) **Youth motor vehicle crash deaths (ages 15 and older)**
- (AHO8) **Adolescent homicide deaths**
- (AHO9) **Adolescent suicide deaths**
- (AHO10) Adolescents with newly diagnosed STD

– *Behavioral Health: Mental Health, Alcohol, Other Drugs & Tobacco*

- (BHO1) % of adults having five or more drinks on an occasion, one or more times in past month; % of high school seniors and young adults 18-24 having five or more drinks on an occasion in the previous two weeks
- (BHO2) % adults who report having 60 or more drinks a month; % of young people who have used alcohol in the past month
- (BHO3) Medical risk for pregnant women—alcohol, tobaccos, and drug use by pregnant women
- (BHO4) Reduce deaths caused by alcohol-related crashes, cirrhosis, and drug-related deaths
- (BHO5) **Reduce suicides; suicide attempts by adolescents; prevalence of mental disorders among adolescents [and] adults living in community**
- (BHO6) Consumers report satisfaction with access, coordination, thoroughness of care, communication, recovery
- (BHO7) % of individuals involved in acute or residential behavioral health treatment program who return to that level of care (or a more intense level) within one year from discharge
- (BHO8) % of individuals completing a formal treatment program who remain involved in aftercare sessions, self-help groups, and have no criminal involvement after one year from program discharge

Notes

¹ See additional measures related to this area of concern in the *Healthy People 2010* section of this workbook.